



# Sinuiju Biota of the Mesozoic in Korea





*The palaeontology research group of the Resources Science Faculty of Kim Il Sung University and an editorial team of the Foreign Languages Publishing House are pleased to publish the picture album Sinuiju Biota of the Mesozoic in Korea with the collection of research findings of palaeontological fossils in the Sinuiju Formation distributed in Paektho-dong and Ryonsang-dong of Sinuiju in the Democratic People's Republic of Korea.*

*Illustrated with various fossils of the Sinuiju Biota, the book shows the history of the evolution and development of terrestrial organisms of the Mesozoic in Korea.*

*The book introduces the Archaeopteryx of Korea and the Ancestral Frog which were named by President Kim Il Sung, the Paekthodong Fossil Reserve in the Sinuiju Biota designated by Chairman Kim Jong Il and fossils of the Paektho Bird named by the respected Comrade Kim Jong Un.*

*It also introduces the geological background, composition and typical fossils of the Sinuiju Biota, new species unearthed in the biota, scientific papers on the biota, Paekthodong Fossil Reserve, study of fossils from the Sinuiju Biota and academic exchanges with regard to it.*

*We have tried to give intuitive and vivid explanations of the Sinuiju Biota of the Mesozoic in Korea with the illustration of photographs of various fossils. We hope that palaeontologists and fossil hobbyists at home and abroad will enjoy this book.*

Won Chol Guk,

Merited Scientist, Professor  
and Doctor, and section  
chief of the Resources  
Science Faculty of  
Kim Il Sung University

## Contents

4	<b>Fossils of Archaeopteryx of Korea and Ancestral Frog</b>
6	<b>Establishment of Paekthodong Fossil Reserve</b>
8	<b>Fossils of Paektho Bird</b>
10	<b>Geological Background of Sinuiju Biota</b>
12	<b>Representative Fossils of Sinuiju Biota</b>
94	<b>New Species Discovered in Sinuiju Biota</b>
100	<b>Scientific Papers on Sinuiju Biota</b>
102	<b>Paekthodong Fossil Reserve</b>
110	<b>Study of Fossils and Academic Exchange</b>







# Sinuiju Biota of the Mesozoic in Korea

About three kilometres east of Sinuiju, North Phyongan Province in the northwestern part of Korea, are Paektho-dong, Ryonsang-dong, Phungso-dong and Thosong-ri. And the Sinuiju Formation is distributed in these areas which embrace the Sinuiju Biota of the Early Cretaceous in the Mesozoic of Korea. The area that exposes the formation is designated as Paekthodong Fossil Reserve, a natural monument, and is under good preservation.

During the study of the Mesozoic fossils that has been conducted in the Paektho-dong area of Sinuiju since the 1960s, typical EEL fauna of the Early Cretaceous in the Mesozoic was confirmed and the Sinuiju Biota was established.

Between late 1980s and early 1990s the fossils of bird, frog and pterosaur were found there.

President Kim Il Sung saw the bird and frog fossils and named them the Archaeopteryx of Korea and Ancestral Frog.

Chairman Kim Jong Il had the Paektho-dong area where the fossils of the Archaeopteryx of Korea were discovered designated as the Paekthodong Fossil Reserve.

Until then, only fragmentary plant fossils were revealed there, so many scientists were engrossed in the study of animal fossils.

It was in 2014 when the palaeontological research team of Kim Il Sung University buckled down to the all-round excavation and investigation of the fossils from the Sinuiju Formation in the Mesozoic

distributed in the areas of Paektho-dong and neighbouring Ryonsang-dong and Phungso-dong.

Accordingly, several thousand pieces of animal and plant fossils, including primordial mammals and dinosaur's tooth, were newly unearthed.

The respected Comrade Kim Jong Un named the fossils of bird with well-preserved quills discovered in the Paektho-dong area the Paektho Bird.

In 2021 palaeontologists gave scientific explanations of the Paektho-dong area of Sinuiju as the birthplace of all groups of vertebrates in the Mesozoic in Korea, and, based on them, established the Sinuiju Biota where a wide variety of animal and plant fossils in the Early Cretaceous are deposited.

The Sinuiju Biota contains the fossils of such vertebrates as fishes, amphibians, reptiles, birds and mammals, the fossils of such invertebrates as insects, conchostracans, bivalves, and gastropods, and several plant fossils.

As there are well-preserved fossils of various insects and the fossils of the Paektho Bird were unearthed and verified recently, the biota has aroused deeper interest at home and abroad.

When the study of the Sinuiju Biota is intensified in the future, it will make a greater stride in the research on the biota of the Early Cretaceous in the Mesozoic of Korea as well as the study of other Mesozoic biota in Northeast Asia.



# Fossils of Archaeopteryx of Korea and Ancestral Frog



## Archaeopteryx of Korea

The fossils of a bird discovered in the Paektho-dong area of Sinuiju, North Phyongan Province, was named the *Archaeopteryx of Korea* on September 27, 1993.



## Ancestral Frog

The frog fossil unearthed in the Paektho-dong area of Sinuiju, North Phyongan Province, was named the *Ancestral Frog* on April 7, 1994.





## Establishment of Paekthodong Fossil Reserve

The Paektho-dong area, where the fossils of Archaeopteryx of Korea were discovered, was designated as the Paekthodong Fossil Reserve on March 19, 1997.




# Fossils of Paektho Bird



0 2 4cm



0 2 4cm



0 2 4cm

## Fossils of Paektho Bird

The fossils of a bird with well-preserved quills unearthed in the Paektho-dong area was named the Paektho Bird on February 13, 2021.



# Geological Background of Sinuiju Biota

The Sinuiju Formation dating to the Mesozoic in Korea is exposed in the areas of Paektho-dong, Ryonsang-dong, Phungso-dong and Thosong-ri in Sinuiju, North Phyongan Province. The formation, mainly distributed in the Sinuiju Basin and structurally located above the older strata (the Lower Proterozoic) along the southwest-northeast direction, is 50km long and 15km wide. It is divided into eight members from bottom to top.

The first member consisting of purple-brown siltstone and well-graded sandstone and mudstone containing invertebrate and plant fossils is deposited unconformably on the Precambrian basement.


The second member consists of andesite and tuffaceous siltstone, where sedimentary layers are interbedded and conformable to the lower member.

The third member is visibly exposed in Paektho-dong, Ryonsang-dong and Phungso-dong and Thosong-ri. And its greyish-green sandstone, grey and black mudstone and siltstone contain a large number of fossils of vertebrates, conchostracans, insects, bivalves, gastropods, ostracods and terrestrial plants. This member is 200-300m thick and rests conformably on the second member, though its thickness varies according to the locations of individual fossils.

The lower part of the fourth member distributed in Paektho-dong consists of sandstone, siltstone, mudstone and slate. Its upper part and the fifth to eighth member consist of andesite, tuffaceous siltstone and sandstone. In the Sinuiju Basin, these strata are not exposed to the surface.

Geologists had previously regarded the geological age of the Sinuiju Formation dating to the Upper Jurassic-Lower Cretaceous.

However, on the basis of insect and bivalve fossils, the age of the third member of the formation was recently reconfirmed as the Lower Cretaceous.





# Representative Fossils of Sinuiju Biota

Most of the fossils in the Sinuiju Biota are found in the third member of the Sinuiju Formation of the Lower Cretaceous and are comprised of invertebrate, vertebrate and plant fossils.

Invertebrate fossils include shellfish, mud snail, Conchostraca, and insects, and vertebrate fossils fish, amphibians (Ancestral Frog), reptiles (tooth of dinosaur and pterosaur), birds (Archaeopteryx of Korea and Paektho Bird), and mammals (primordial animals). And plant fossils contain pteridophytes, *Pteridospermopsida*, horsetails, *Cycadopsida*, and Ginkgoaceae.

Of the fossils produced by the Sinuiju Biota are those of insects, including Odonata, Neuroptera, Orthoptera, Coleoptera, Homoptera, Blattulidae, Hymenoptera, and Hemiptera.



Reconstruction map of the natural environment of the Sinuiju Biota





Representative Fossils of Sinuiju Biota

# Gastropods

Gastropods are found generally together with bivalves in the either greyish-black siltstone or greyish-black silty mudstone. They are limited in terms of genus and species and badly preserved as compared to other fossils so far unearthed in the Sinuiju Biota.



PS2014-12



PS2015-3



PS2015-04



PS2020-19



PS2016-25



PS2016-56



PS2014-22



PS2014-41



PS2015-09



PS2014-13



PS2014-65



PS2018-27



PS2016-08



PS2014-19



PS2016-44



PS2014-20



PS2015-14



PS2015-16



Representative Fossils of Sinuiju Biota

# Bivalves

Bivalves are found both in the second and third members of the Sinuiju Formation, and in fairly concentrated way in certain horizons of the third member.

Such bivalves as *Sphaerium anderssoni*, *Arguniella lingyuanensis*, *Arguniella yanshanensis*, and *Ferganoconcha dapingshanensis* have been discovered in the grey siltstone and greyish-black silty mudstone of the formation, and they preserve their shell forms and structures in a vivid way.



PS2020-06



PS2020-13



PS2016-09



PS2018-11





PS2015-07



PS2016-17



PS2016-29



PS2016-36



PS2015-02



PS2015-11



PS2015-11



PS2020-21



PS2021-12



PS2020-08



PS2015-13



PS2014-04



PS2015-20



PS2021-20



PS2020-11



PS2015-19



PS2018-02

PS2022-09





PS2020-14



PS2015-23



PS2021-07

PS2021-03



PS2015-15



PS2016-20





PS2021-10



PS2016-21



PS2015-31



PS2018-17



PS2021-14



Representative Fossils of Sinuiju Biota

# Conchostracans

The Sinuiju Biota contains conchostracans generally in the layers of the third member of the Sinuiju Formation, and in many cases they occur with insects and plants.

The Paektho-dong and Ryonsang-dong areas produced various species of conchostracans, including *Yanjiestheria yanjiensis*, *Yanjiestheria dalaziensis*, *Eosestheria middendorffii*, *Eosestheria ovata*, and *Longjiangestheria cericula*, most of them belonging to *Yanjiestheria* and *Eosestheria*.

They are found in the yellowish-brown and grey mudstone, and a rich assemblage of well-preserved fossils in several layers.



PS2014-06



PS2015-33




PS2018-08

PS2020-20







PS2015-34



PS2015-34



PS2022-10



PS2016-23



PS2015-12



PS2022-27



PS2020-31

PS2014-14



PS2020-22



PS2018-08



PS2018-08



PS2018-33



PS2016-11





PS2015-03



PS2015-24



PS2021-11



PS2021-25



PS2020-23



PS2020-23



PS2021-19



PS2014-24



PS2018-06



PS2018-12



PS2021-18



PS2021-30

PS2022-13



PS2022-21





PS2020-18



PS2018-25



PS2016-40



PS2020-36



PS2021-13



PS2016-22



PS2021-29



PS2020-17



PS2018-26



PS2020-33



PS2020-37





PS2014-15



PS2015-29



PS2015-41



PS2016-37



PS2015-17



PS2015-17



PS2017-01



PS2021-21



PS2016-35



PS2016-24



PS2016-43



PS2016-55



PS2015-42






PS2021-15



PS2022-20



PS2022-22



PS2016-53



PS2020-40



PS2016-61



PS2021-04



PS2016-64




PS2022-23



PS2022-02



PS2022-07



PS2021-16



PS2022-06



PS2016-63



PS2015-50



PS2014-21



PS2015-51



PS2014-27



PS2018-30



# Representative Fossils of Sinuiju Biota

## Insects

Insects are the most diversified invertebrate group of the Sinuiju Biota.

Their fossils are rich in the greyish-green and grey mudstone and silty mudstone in the third member of the Sinuiju Formation.

Here *Ephemeropsis trisetalis* Eichwald, 1864 (Ephemeroptera), is found in all areas of the third member of the Sinuiju Formation, and they are well preserved and abundant.

The Sinuiju Biota also contains such insects as Odonata, Neuroptera, Orthoptera, Coleoptera, Homoptera, Blattulidae, Hymenoptera, and Hemiptera.

Among the new species are *Angarosphex baektoensis* Jon et al., 2019, *Sinuijuhelorus baektoensis* Jon et al., 2019, *Sinuijus baektoensis* Fang et al., 2019, *Habroblattula sinuijunensis* Fang et al., 2019, *Sinuijularis baektoensis* Jon et al., 2019, *Karataus ryonsangensis* So et Won, 2021, *Khasurtella ryonsangi* So et Won, 2021, *Pompilopterus ryonsangensis* So et Won, 2021, *Pompilopterus ryonsangensis* So et Won., 2021, *Stenophlebia ryonsangensis* Won et al., 2021, *Stellularis sinuijuensis* Jon et al., 2020, *Sinuijumantispa ryonsangia* Won et So, 2021, *Sinuijupodagrion ryonsangae* So et Won, 2021, *Aenigmoilus ryonsangensis* So et Won, 2020, and *Stellularis ryonsangensis* So et Won, 2021.

Insect fossils are found simultaneously with conchostracan and plant fossils.



PS2015-01



PS2016-12



PS2019-07




PS2014-11

PS2020-34







PS2020-41



PS2015-10



PS2016-62




PS2018-31



PS2021-26



PS2020-51



PS2020-10





PS2021-32




PS2020-45

PS2021-34



PS2021-34








PS2018-03



PS2020-50



PS2020-39



PS2016-13



PS2021-34



PS2020-42





PS2020-52



PS2015-21



PS-2015-25



PS2021-35



PS2021-36



PS2018-32



PS2018-16

PS2020-53



PS2015-60



PS2020-67



PS2018-01



PS2020-66



PS2015-05



PS2020-68

PS2014-26





PS2021-37



PS2018-04

PS2020-02



PS2020-69



PS2020-04





PS2014-02



PS2022-11



PS2022-17



PS2018-29



PS2020-70

PS2021-38



PS2019-10



PS2019-13



PS2015-35



PS2018-13



PS2022-26







PS2018-34



PS2018-14



PS2018-10



PS2015-49



PS2019-01



PS2019-09



PS2016-65



PS2014-23



PS2015-52



PS2019-11





PS2015-53



PS2016-66



PS2015-54



PS2021-39



PS2015-61



PS2018-31



PS2021-44



PS2018-35





PS2021-40



PS2021-50



PS2019-12



PS2016-67



PS2020-71



PS2018-36



PS2016-10



PS2016-30



PS2021-41



PS2014-25



PS2016-52



PS2021-42



PS2019-14



PS2021-45





PS2019-15



PS2020-72



PS2022-18



PS2018-37



PS2019-16



PS2014-28



PS2021-46



PS2021-55



PS2018-38



PS2019-17



PS2019-18





PS2014-29



PS2021-47



PS2019-19



PS2016-54



PS2018-40



PS2020-73



PS2018-39



PS2021-48



PS2022-24



PS2021-49



PS2020-74





PS2019-20



PS2018-41



PS2022-25



PS2019-21



PS2022-30



PS2022-28



PS2019-22



PS2016-68



PS2021-51



PS2018-42



PS2021-01



PS2021-52



PS2018-43



PS2022-29



PS2021-53





PS2021-02



PS2021-03



PS2018-44



PS2019-23



PS2014-30



PS2018-45



PS2015-62



PS2019-24



PS2020-75



PS2021-54



PS2021-08





PS2021-55



PS2016-69



PS2015-63

PS2019-25



PS2020-76



PS2019-26



PS2022-31







PS2019-27




PS2021-56



PS2022-32



PS2018-47



PS2020-100



PS2020-110



PS2019-09



PS2019-28



PS2018-46



PS2020-77





PS2020-78



PS2014-31



PS2018-48



PS2021-57



PS2020-79



PS2022-33

PS2021-58



PS2015-64



PS2022-34

PS2018-49



PS2019-29





Representative Fossils of Sinuiju Biota

Fish

PS2015-65

Fishes are often discovered in the greyish-black silty mudstone of the third member of the Sinuiju Formation, and rarely in the greyish-green mudstone.

*Lycopera davidii*, *Lycopera muroii* and

*Peipiaosteus* sp. 1, *Peipiaosteus* sp. 2 are included in the fossils from the Sinuiju Biota.


*Lycopera* are the most common species in the Sinuiju Biota, and they preserve shapes and

structures fairly well.

In the lower layer of the member containing fish fossils are also produced remains of bivalves, gastropods and plants, while in the upper layer insects, conchocratans and plants.







PS2015-66



PS2014-32



PS2016-70



PS2021-59



PS2016-71




PS2015-67

PS2014-33

PS2021-60







PS2014-34

PS2015-68



PS2016-72

PS2015-69



PS2016-73





PS2019-30



PS2015-70



PS2016-74



PS2018-50



PS2016-01



PS2016-02



PS2014-37

PS2019-31

PS2019-32



PS2014-35



PS2014-36



PS2015-71



PS2020-80



PS2018-51



PS2015-72







PS2014-38



PS2014-03



PS2014-05



PS2014-07



PS2021-61



PS2015-05



PS2016-03



PS2019-33



Representative Fossils of Sinuiju Biota

# Frog

In the early 1990s a frog fossil was discovered in the Sinuiju Biota. Found in the greyish-green silty mudstone of the third member of the Sinuiju Formation distributed in Paektho-dong, the fossil has most of its skeletons, except its head, that are relatively well-preserved. It has similar morphological features with *Liaobatrachus grabaui* Ji and Ji, 1998. The fossil is named the Ancestral Frog of Korea.



PS200121

Representative Fossils of Sinuiju Biota

# Pterosaurs

It is known that the Sinuiju Biota contains two pterosaur categories, all unearthed in the third member of the Sinuiju Formation.

The fossils prove that pterosaurs existed in the Paektho-dong area in the Early Cretaceous in the Mesozoic.

The fossil of the pterosaur (PS198922) is thought to belong *Jeholopterus ningchengensis* Wang et al., 2002.



PS198922



PS199921

Representative Fossils of Sinuiju Biota

# Dinosaur's Tooth

The fossilized dinosaur's tooth was found in the third member of the Sinuiju Formation distributed in Paektho-dong in 2016.

The tooth, smooth-surfaced and a little crooked backward, has close sawtoothed denticles along the anterior and posterior margins.



PS201643



Representative Fossils of Sinuiju Biota

# Birds

A bird fossil that was first unearthed in the early 1990s in the grey silty mudstone layer of the third member of the Sinuiju Formation preserves some parts of the skull and toe bones. The fossilized bird is named the Archaeopteryx of Korea.

And another fossil unearthed in the Paektho-dong area of the Sinuiju Biota in 2020 has well-preserved skull, toe bones and eleven primary feathers of a bird, which is named the Paektho Bird.

Both bird fossils are registered as the DPRK’s natural monuments.



PS201403



PS2020723

Representative Fossils of Sinuiju Biota

# Mammal

The mammalian fossil has been recently discovered in the Sinuiju Biota.

Though it has no skull and forelimbs, the fossil preserves relatively clear structures of hindlimbs and chest. Judging from the characteristics of its well-preserved toe bones, it belongs to the Cretaceous mammal.



PS201641



Representative Fossils of Sinuiju Biota  
Plants

Plants are found alike in the second, third and fourth members of the Sinuiju Formation, but most of them are in the third member.

Especially, fossils of Pteridophytes, *Pteridospermopsida*, *Cycadopsida*, Ginkgoaceae, Sphenophytes, and Conifers produced in the areas of Ryonsang-dong and Paektho-dong are in rather good preservation, but they are not so varied in terms of species and genus.

Most common plants recently unearthed in the third member of the Sinuiju Formation include *Coniopteris ermolaevii*, *Coniopteris vachrameevii*, *Equisetum* sp., *Czekanowskia rigida*, and *Baiera borealis*.



PS2022-35



PS2016-05



PS2016-75



PS2015-80



PS2015-81



PS2022-36





PS2015-73



PS2015-82



PS2016-06



PS2020-81



PS2022-37



PS2021-62



PS2016-07



PS2015-74



PS2016-100



PS2015-83



PS2014-08



PS2016-101





PS2021-63



PS2021-64



PS2020-82



PS2014-39



PS2016-102



PS2016-80



PS2021-64



PS2016-81



PS2016-103



PS2014-40



PS2014-42



PS2021-65

PS2016-105







PS2021-66



PS2022-51



PS2022-55



PS2022-60



PS2020-83



PS2019-35



PS2022-58

PS2022-50

The fossils from the Sinuiju Biota contain a variety of invertebrates of the Early Cretaceous in the Mesozoic, including bivalves, gastropods, conchostracans, and insects, and major five groups of vertebrates—fishes, amphibians, reptiles, birds and mammals.

The palaeontological circle of the DPRK named the biota the Sinuiju Biota, which contains a wide variety of fossils of lake-dwelling animals and plants of the Early Cretaceous.

The establishment of the Sinuiju Biota is of great academic significance in explaining the evolution and development of terrestrial ecosystem and living creatures in the Mesozoic of Korea.




# New Species Discovered in Sinuiju Biota


Palaeontologists of the DPRK newly discovered and verified many insect fossils through their study of the Sinuiju Biota.

Below are illustrated insect fossils of new species.

- A-*Karataus ryonsangensis* Won et al., 2021
- B-*Khasurtella ryonsangi* Won et al., 2021
- C-*Pompilopterus ryonsangensis* Won et al., 2021
- D-*Pompilopterus ryonsangensis* Won et al., 2021




*Stenophlebia ryonsangensis* Won et al., 2021




*Stellularis sinuijuensis* Jon et al., 2020






**B**

*Sinuijumantispa ryonsangia* Won et So, 2021




**C**

*Angarosphex baektoensus* Jon et al., 2019



**B**



**D**

*Sinuijupodagrion ryonsangae* Won et So, 2021









*Aenigmoilus ryonsangensis* So et Won, 2020



*Stellularis ryonsangensis* So et Won, 2021



*Sinuijus baektoensis* Fang et So, 2020



*Sinuijupodagrion ryonsangae* So et Won, 2021



# Scientific Papers on Sinuiju Biota

Research findings of the Sinuiju Biota were published in nearly 20 scientific papers and international journals, such as *Cretaceous Research*, *Palaeoworld*, and *Palaeontological Journal*.

It consists of greyish-green sandstones, grey to black mudstones and siltstones and yields vertebrate fossils, insects, conchostracans, bivalves, gastropods, ostracodes, and terrestrial

History of Paleontology of Sinuiju Biota follows

Sinuiju City is located in the southwestern part of the Sinuiju basin, which is a tuffaceous siltstone overlying the Cretaceous and Paleogene basement.



**Short communication**  
**The first Orthoptera (Insecta) from the Lower Cretaceous of Democratic People's Republic of Korea**  
 KwangSik So <sup>a</sup>, CholGuk Won <sup>a</sup>, KumSik Han <sup>a</sup>, Chunpeng Xu <sup>b,c</sup>, Bo Wang <sup>b</sup>, Yan Fang <sup>b,\*</sup>

<sup>a</sup> Department of Paleontology, Faculty of Geology, Kim Il Sung University, Pyongyang, Democratic People's Republic of Korea  
<sup>b</sup> State Key Laboratory of Palaeontology and Stratigraphy, Nanjing Institute of Geology and Palaeontology and Center for Excellence in Life and Palaeoenvironment, Chinese Academy of Sciences, Nanjing, 210008, China  
<sup>c</sup> University of Science and Technology of China, Hefei, 230026, China



**Short communication**  
**New Mesozoic insect fossils from the Democratic People's Republic of Korea**  
 SuHyang Jon <sup>a</sup>, CholGuk Won <sup>a</sup>, KwangSik So <sup>a,\*</sup>, TuYong Nam <sup>b</sup>

<sup>a</sup> Department of Paleontology, Faculty of Geology, Kim Il Sung University, Pyongyang, Democratic People's Republic of Korea  
<sup>b</sup> Institute of Zoology, South Academy of Science (SAS), Pyongyang, Democratic People's Republic of Korea

## ARTICLE INFO

**Article history:**  
 Received 4 October 2018  
 Received in revised form 8 January 2019  
 Accepted in revised form 18 February 2019  
 Available online 22 February 2019

**Keywords:**  
 Mesozoic  
 Sinuiju Formation

## ABSTRACT

The insect fossils were collected from the Lower Cretaceous Sinuiju Formation in Paekto-dong, Sinuiju City, North Pyongan Province, Democratic People's Republic of Korea (DPRK). A new genus with a new species, *Sinuijuhirsutus bakosensis* gen. et sp. nov., is described and assigned to the family Heloridae. In addition, a new species, *Angarosphex bakosensis* sp. nov. (Angarospheridae) is also described. The new species not only indicate that the genus *Angarosphex* reached the DPRK in the Early Cretaceous, but also broaden the distribution and diversity of the family Heloridae.

© 2019 Elsevier Ltd. All rights reserved.



**New Evidence for a Cretaceous Age for a Mesozoic Nanoscale Brachiopod from Paekto-dong, Sinuiju City, The Democratic People's Republic of Korea**

CholGuk WON, KwangSik SO<sup>a</sup>, and SuHyang JON<sup>a</sup>

<sup>a</sup> Department of Paleontology, Faculty of Geology, Kim Il Sung University, Pyongyang, Democratic People's Republic of Korea

**Abstract:** The Sinuiju Formation in Paekto-dong, Sinuiju City in the Democratic People's Republic of Korea has yielded numerous brachiopod fossils, which is the first occurrence of such in the DPRK. Based on these fossil specimens, a new Cretaceous brachiopod, *Angarosphex bakosensis*, the youngest Cretaceous brachiopod, is named. The morphology includes *Angarosphex bakosensis*, *A. bakosensis* and *Angarosphex bakosensis* and can be compared with the fossil brachiopod. The age of the Sinuiju Formation is also identified on the basis of the location and the presence of a brachiopod (*Angarosphex bakosensis*) of the Cretaceous. The Sinuiju Formation is not typical for the Lower Cretaceous in the DPRK.

Keywords: Geology, Paleontology, Sinuiju Formation, Early Cretaceous, DPRK



**A new Odonatan (Stenophlebiidae) from the Lower Cretaceous Sinuiju Formation of North Pyongan Province, the Democratic People's Republic of Korea**


Chol-Guk Won, Kwang-Sik So<sup>a</sup>, Hui-Chol Kim

<sup>a</sup> Department of Paleontology, Faculty of Geology, Kim Il Sung University, Pyongyang, Democratic People's Republic of Korea

Received 4 October 2018; received in revised form 23 December 2018; accepted 1 February 2019

## Abstract

A new Cretaceous Odonatan (Stenophlebiidae) from the Lower Cretaceous Sinuiju Formation in Paekto-dong, Sinuiju City, North Pyongan Province, Democratic People's Republic of Korea (DPRK). A new genus with a new species, *Sinuijuhirsutus bakosensis* gen. et sp. nov., is described and assigned to the family Heloridae. In addition, a new species, *Angarosphex bakosensis* sp. nov. (Angarospheridae) is also described. The new species not only indicate that the genus *Angarosphex* reached the DPRK in the Early Cretaceous, but also broaden the distribution and diversity of the family Heloridae.



## Short communication

**A new species of Proceropidae (Insecta: Hemiptera) from the Lower Cretaceous of Democratic People's Republic of Korea**

Suhyang Jon<sup>a</sup>, KwangSik So<sup>a</sup>, CholGuk Won<sup>a</sup>, LiJong Hyon<sup>a</sup>

<sup>a</sup> Department of Paleontology, Faculty of Geology, Kim Il Sung University, Pyongyang, Democratic People's Republic of Korea

## ARTICLE INFO

**Article history:**  
 Received 12 March 2018  
 Received in revised form 8 January 2019  
 Accepted in revised form 18 February 2019  
 Available online 22 February 2019

**Keywords:**  
 Mesozoic  
 Sinuiju Formation

## ABSTRACT

The insect fossils were collected from the Lower Cretaceous Sinuiju Formation in Paekto-dong, Sinuiju City, North Pyongan Province, Democratic People's Republic of Korea (DPRK). A new genus with a new species, *Sinuijuhirsutus bakosensis* gen. et sp. nov., is described and assigned to the family Heloridae. In addition, a new species, *Angarosphex bakosensis* sp. nov. (Angarospheridae) is also described. The new species not only indicate that the genus *Angarosphex* reached the DPRK in the Early Cretaceous, but also broaden the distribution and diversity of the family Heloridae.

© 2019 Elsevier Ltd. All rights reserved.

## 1. Introduction

The insect fossils were collected from the Lower Cretaceous Sinuiju Formation in Paekto-dong, Sinuiju City, North Pyongan Province, Democratic People's Republic of Korea (DPRK). A new genus with a new species, *Sinuijuhirsutus bakosensis* gen. et sp. nov., is described and assigned to the family Heloridae. In addition, a new species, *Angarosphex bakosensis* sp. nov. (Angarospheridae) is also described. The new species not only indicate that the genus *Angarosphex* reached the DPRK in the Early Cretaceous, but also broaden the distribution and diversity of the family Heloridae.



Fig. 1. Geographic map showing the Sinuiju locality.



**Short communication**  
**First record of a cockroach (Insecta: Blattaria: Mesoblattinidae) from the Sinuiju Formation of the Democratic People's Republic of Korea**

Kwang-Sik So<sup>a</sup>, Chol-Guk Won, Chol-Jun Ri, Su-Hyang Jon

<sup>a</sup> Department of Paleontology, Faculty of Geology, Kim Il Sung University, Pyongyang, Democratic People's Republic of Korea

## ARTICLE INFO

**Article history:**  
 Received 3 April 2020  
 Received in revised form 19 November 2020  
 Accepted in revised form 19 November 2020  
 Available online 19 November 2020

**Keywords:**  
 DPRK  
 Sinuiju Formation  
 Yulian Formation  
 Paekto-dong  
 Mesoblattinidae

## ABSTRACT



**Short communication**  
**First record of a beetle (Coleoptera: Cupedidae) from the Lower Cretaceous of the Democratic People's Republic of Korea**

CholGuk Won, SuHyang Jon, KwangSik So<sup>a</sup>, LiJong Hyon

<sup>a</sup> Department of Paleontology, Faculty of Geology, Kim Il Sung University, Pyongyang, Democratic People's Republic of Korea

## ARTICLE INFO

**Article history:**  
 Received 15 March 2019  
 Received in revised form 20 August 2019  
 Accepted in revised form 13 October 2019  
 Available online 25 October 2019

**Keywords:**  
 DPRK  
 Sinuiju Formation  
 Cretaceous  
 Cupedidae

## ABSTRACT

The beetle species *Zygadenia lusi* (Coleoptera: Cupedidae: Ommatidae) is found in the materials collected in the Sinuiju Formation from Paekto-dong, Sinuiju City, North Pyongan Province, Democratic People's Republic of Korea (DPRK). This discovery extends the geographic distribution of *Zygadenia* in the Mesozoic.

© 2019 Elsevier Ltd. All rights reserved.

## 1. Introduction

Paleoentomologists considered the genera related to *Omma* with conjoined pronotum as the subfamily Ommatidae of the family Cupedidae (Ponomarenko, 1965; Krasnoluk et al., 2016; Ponomarenko et al., 2019; etc.) because of a small hump of this group from other cupedids. The extant Ommatidae is a very small group including two genera: *Omma* Newman, 1839 and *Tetraphaleria* Waterhouse, 1901 with six species (Sharp et al., 1912), and their distribution is only restricted to the tropical belt of the southern hemisphere. However, it was abundant in the Mesozoic, and many Jurassic and Cretaceous fossil ommatids from China, Siberia, Central Asia and western Europe have been reported by Ponomarenko (1965, 1997, 2000), Yan et al. (2005), Sieroski and Dabkowski (2008), Jurekiewicz et al. (2017) and Krasnoluk et al. (2016). The beetles belonging to the extinct tribe Notocupedini (Ponomarenko, 1966) (Coleoptera: Cupedidae: Ommatidae) are widespread in Mesozoic deposits of eastern Pangaea (Jurekiewicz et al., 2015).

Recently we recovered beetles belonging to the extinct tribe Notocupedini (Coleoptera: Cupedidae: Ommatidae) from the Sinuiju Formation near Paekto-dong, Sinuiju City, North Pyongan Province.

\* Corresponding author.  
 E-mail address: yu.suhyang@pyongyang.ac.kr (K. So).

https://doi.org/10.1016/j.cretres.2019.104636  
 0195-6671/© 2019 Elsevier Ltd. All rights reserved.

The Sinuiju Formation is mainly of lacustrine strata intercalated with volcanics (Park and Kim, 1996). It contains a large number of fossils, including vertebrate fossils, insects, conchostracans, bivalves, gastropods, ostracodes, and terrestrial plant fossils (Park and Kim, 1996). The age of the Sinuiju Formation is Early Cretaceous (Yan et al., 2010). Specimen was examined dry and wetted with ethanol under a Leica S6D digital camera. Photographs were taken using a Leica S6D digital camera and line drawings were readjusted on photographs using image-editing software CorelDraw. The specimen examined is deposited in the Laboratory of Paleontology, Faculty of Geology, Kim Il Sung University.



Fig. 1. Geographic map and outcrop showing the Sinuiju locality.





# Paekthodong

## Fossil Reserve



The Paektho-dong area that contains various animal and plant fossils of the Early Cretaceous has been designated as the Paekthodong Fossil Reserve, a natural monument, in 1997.

The reserve is distributed in four separate sections.

The reserve has fair-arranged geological cross sections and well-preserved fossils in various kinds, so it is of importance in the study of the Sinuiju Biota of the Lower Cretaceous in the Mesozoic.

### Section 1

Major fossils produced in Section 1 include fishes, bivalves, gastropods, conifers and pteridophytes deposited in the greyish-black silty shale and greyish-green silty mudstone.

Fish fossils were discovered in abundance there.



## Section 2

The fossils of the Paektho Bird was unearthed here, together with a large number of insect and conchostracan fossils and some plant fossils. The most representative insect fossil is *Ephemeropsis trisetalis* Eichwald, 1864. The layer is composed of greyish-green silty mudstone, yellowish-brown siltstone and greyish-green sandstone.








## Section 3

Section 3 produces mainly insect, conchostracan, bivalve and gastropod fossils, and consists of greyish-green silty mudstone, greyish-green sandstone and yellowish-brown sandstone.







## Section 4

The fossils of the Archaeopteryx of Korea, pterosaur, fishes, bivalves, gastropods, pteridophytes, and Ginkgoaceae were discovered in the section. The section consists of greyish-black sandstone, siltstone and silty mudstone.





# Study of Fossils and Academic Exchange

Excavation and Study of Fossils from Sinuiju Biota in Paektho-dong and Ryonsang-dong



The DPRK's palaeontologists intensified the study of fossils from the Sinuiju Formation since 2014, and unearthed and verified a lot of fauna and flora fossils in Paektho-dong and Ryonsang-dong of Sinuiju.



# Study of Fossils and Academic Exchange

## Academic Discussions with Chinese palaeontologists

Fossil samples from the Sinuiju Biota of the Mesozoic are exhibited in the palaeontological hall of the Natural History Museum built at the foot of Mt Taesong, DPRK.


Professor Sun Ge, director of the Paleontological Institute of the Shenyang Normal University and also head of the Paleontological Museum of Liaoning in China, and other Chinese palaeontologists visited the Natural History Museum and looked round its palaeontological hall.

Palaeontologists of the DPRK and China exchanged their scientific opinions on the evolution and development of the Mesozoic organisms.

Academic discussion with the palaeontologists of the Shenyang Normal University of China at the palaeontological hall of the Natural History Museum







The DPRK palaeontologists and those of the Shenyang Normal University of China agreed to intensify their academic exchange on the Mesozoic biota in the DPRK and northeast China.



Academic discussion with the palaeontologists of the Shenyang Normal University of China



Joint research and academic seminar with the Nanjing Institute of Geology and Palaeontology of Chinese Academy of Sciences

The DPRK palaeontologists met with Prof. Zhan Renbin, director of the Nanjing Institute of Geology and Palaeontology of Chinese Academy of Sciences, and Prof. Wang Bo. They discussed the orientation of their joint research and collaborated in the research and compilation of scientific papers on insect fossils.







## Study of Fossils and Academic Exchange

### Academic Exchange with Palaeontologists from Mongolian Academy of Sciences

Mongolian palaeontologists headed by Prof. Khishigjav Tsogtbaatar, director of the Institute of Palaeontology and Geology of the Mongolian Academy of Sciences, visited Kim Il Sung University and agreed to conduct joint study and academic exchange with the DPRK counterparts.



Mongolian palaeontologists visiting the DPRK



## Study of Fossils and Academic Exchange

### International Seminars

The DPRK palaeontologists participated in several international meetings, including the International Symposium on Cretaceous Biota and K-Pg Boundary in Jiayin of Heilongjiang and Jiayin Forum on Fossil Protection, China, the Fourth International Symposium on Geosciences in Northeastern Asia, and the Fourth Council of the International Centre of Geosciences Research and Education in Northeastern Asia. The meetings discussed the issues on the joint study of the palaeontological evolution and development in the Mesozoic of Korea, and scientific papers were read.



International Symposium on Cretaceous Biota and K-Pg Boundary in Jiayin of Heilongjiang, China







Many DPRK and foreign scientists are paying great attention to the Sinuiju Biota of the Mesozoic in Korea.  
The Sinuiju Biota will further produce new and valuable fossils in larger numbers and thus accelerate the related academic exchanges.



Fourth International Symposium on Geosciences in Northeast Asia

Fourth Council of the International Centre of Geosciences Research and Education in Northeast Asia





Editor: Kim Kuk Chol

Author: Won Chol Guk, Merited Scientist, Professor and Doctor, section chief of  
Resources Science Faculty of Kim Il Sung University

Photo: Pyon Chan U, Song Tae Hyok, Yun Hyok

## **Sinuiju Biota of the Mesozoic in Korea**

---

Published by the Foreign Languages Publishing House  
Printed at the Printing Plant of the Foreign Languages  
Publishing House  
Issued in December Juche 111 (2022)

---







Foreign Languages Publishing House, DPRK  
Juche 111 (2022)